Advancing Sustainable Agriculture:

The Role of EFB Biochar in Fertilizer Reduction



Organics

Advancing Sustainable Agriculture: The Role of EFB Biochar in Fertiliser Reduction

Sustainable agricultural practices are at the forefront of addressing the growing global concern over environmental degradation, excessive use of chemical fertilisers, and the depletion of natural resources. One innovative solution gaining traction is the use of biochar, particularly biochar derived from empty fruit bunches (EFB). This sustainable approach not only reduces the dependency on chemical fertilisers but also enhances soil fertility, promotes environmental benefits, and supports long-term agricultural productivity.

Pyrolysis of EFB

Understanding Biochar and its Production

Biochar is a carbon-rich material produced through the pyrolysis of organic biomass. Its unique structure, formed under high temperature conditions, makes it highly effective in improving soil properties. Biochar has been widely recognised for its ability to enhance soil fertility by increasing nutrient availability, enhancing microbial activity, improving water retention, and reducing soil bulk density.

Empty Fruit Bunch (EFB) biochar is produced specifically from the agricultural waste of oil palm cultivation. The pyrolysis process involves heating the EFB under controlled conditions, producing a material rich in essential nutrients like potassium, phosphorus, and carbon, which are crucial for plant growth. EFB biochar is produced through different pyrolysis techniques, such as conventional pyrolysis and microwave-assisted pyrolysis, with each method optimising the carbon content and nutrient levels of the resulting biochar.

Biochar's Benefits for Soil Health and Fertility

One of the most significant benefits of biochar application is its ability to improve soil fertility. The porous nature of biochar enhances soil structure, making it easier for plants to access nutrients and water. In particular, EFB biochar improves soil health to make vital nutrients like phosphorus and potassium more accessible, and it is highly effective at retaining nitrogen in the soil, preventing nutrient loss and ensuring it is available for healthy plant growth.

Research has shown that co-applying EFB biochar with chemical fertilisers can significantly improve soil nutrient profiles, particularly in rice cultivation, by enhancing nutrient retention and reducing nutrient leaching.



Process flow diagram indicating how biochar is produced

Biochar's ability to reduce nutrient leaching is one of its most notable advantages. In regions where intensive farming practices are common, nutrient leaching is a significant concern, contributing to water pollution and soil degradation. Studies have demonstrated that biochar can reduce the leaching of key nutrients such as nitrogen and phosphorus, ensuring better nutrient retention in the soil. This benefit not only improves soil health but also contributes to environmental sustainability by minimising nutrient runoff into water bodies.

Moreover, biochar's capacity to stimulate microbial activity is an important factor in improving soil health. By creating an ideal environment for soil microorganisms, biochar promotes nutrient cycling and enhances the soil’s organic carbon content, both of which contribute to long-term soil fertility.

Empty Fruit bunch

Biochar produced from EFB

Environmental Impact and Greenhouse Gas Mitigation

The environmental implications of using EFB biochar are significant. Beyond improving soil health, biochar also contributes to mitigating climate change. The stable carbon content of biochar means it can sequester carbon in the soil for hundreds or even thousands of years, providing a long-term method of carbon storage. This feature is particularly beneficial in the context of rising greenhouse gas emissions.

EFB biochar has also been shown to reduce methane (CH4) emissions, a potent greenhouse gas. Research has highlighted that biochar can significantly lower methane emissions in rice paddies by up to 43%. This property positions biochar as a promising tool for addressing climate change while improving agricultural productivity.

Reducing Chemical Fertiliser Use

The co-application of biochar with chemical fertilisers is a sustainable practice that reduces the need for synthetic fertilisers, which are often expensive, energy-intensive, and environmentally harmful. The incorporation of biochar into soil enhances nutrient availability, which means that farmers can use fewer chemical fertilisers without sacrificing crop yields. This reduction in fertiliser use not only lowers farming costs but also diminishes the environmental impact of fertiliser overuse, such as soil degradation, water pollution, and greenhouse gas emissions.

In regions like Southeast Asia, where intensive cultivation practices often lead to soil depletion, the use of EFB biochar offers a practical solution to enhance productivity while minimising environmental harm. Studies have demonstrated that EFB biochar can help sustain crop yields while reducing the reliance on conventional fertilisers, particularly in crops like rice, which are heavily dependent on fertiliser inputs.

Summary of the uses of biochar from EFB

Economic Implications for Farmers

From an economic perspective, the use of biochar offers substantial benefits. By reducing the need for chemical fertilisers, farmers can save on input costs. In addition, the increased availability of nutrients in the soil leads to improved crop yields. For example, studies have shown that the combined use of EFB biochar and chemical fertilisers can result in yield improvements of over 36%, particularly during the critical early growth stages of rice. These enhanced yields translate into higher revenue for farmers, offsetting the initial investment required for biochar production and application.

Moreover, biochar's role in improving long-term soil health means that farmers can enjoy sustained productivity with reduced input costs over time. Healthier soils with improved nutrient cycling and water retention capacities contribute to more resilient crops, reducing the likelihood of crop failure and the need for additional agricultural inputs.



Yearly Fertiliser Costs (USD) for a 10,000-ha Palm Plantation



Fertiliser Costs for a Palm Plantation

Challenges and Barriers to Widespread Adoption

While the potential benefits of EFB biochar are clear, several challenges remain in its widespread adoption. One significant barrier is the economic cost associated with biochar production and application. The initial investment required for biochar production, equipment, and labour can be prohibitive for smallholder farmers, especially in developing regions. Although the long-term savings from reduced fertiliser use can offset these initial costs, many farmers may be reluctant to invest in biochar without financial support or incentives.

Another challenge lies in the variability of biochar's effectiveness depending on soil type, climate conditions, and agricultural practices. Research has shown that while biochar can improve soil properties in certain contexts, its effectiveness is not universally applicable. Therefore, it is crucial for farmers to conduct site-specific trials and adapt biochar applications to their unique environmental conditions.

Additionally, there is a need for greater awareness and education among farmers about the benefits and proper application of biochar. Many farmers may not be familiar with biochar technology or its potential to improve soil fertility and reduce fertiliser use. Extension services and educational programs are vital to bridging this knowledge gap and encouraging the adoption of biochar-based farming practices.

Policy Support and Future Directions

To overcome these barriers, targeted policies and incentives are essential. Governments and agricultural organisations can support the use of biochar by providing financial incentives, such as subsidies or grants, to encourage its adoption. Additionally, clearer regulatory frameworks and guidelines on biochar application would help farmers navigate the implementation process and ensure its effective use. Further research and development are also necessary to optimise biochar production methods, improve its cost-effectiveness, and refine its applications for different agricultural systems. By addressing the economic, environmental, and knowledge

barriers, biochar could play a key role in advancing sustainable agricultural practices and reducing the reliance on chemical fertilisers.

Conclusion

EFB biochar offers a promising solution for reducing fertiliser use, enhancing soil fertility, and promoting sustainable agricultural practices. Its ability to improve soil health, reduce nutrient leaching, and mitigate greenhouse gas emissions makes it a valuable tool in the fight against environmental degradation. With continued research, policy support, and farmer education, biochar has the potential to revolutionise agricultural practices, providing long-term benefits for both farmers and the environment.